

EXTREMELY RED OBJECTS: AN X-RAY DICHOTOMY

M. BRUSA^{1,2}, A. COMASTRI², E. DADDI³, A. CIMATTI⁴, M. MIGNOLI², L. POZZETTI²¹ Dipartimento di Astronomia, Università di Bologna, via Ranzani 1, I-40127 Bologna, Italy² INAF – Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italy³ European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching, Germany⁴ INAF – Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-55025 Firenze, Italy

Draft version February 5, 2008

ABSTRACT

We analyze the X-ray properties of a near-infrared selected ($K_s < 20$) sample of spectroscopically identified Extremely Red Objects ($R - K_s > 5$) in a region of the *Chandra* Deep Field–South using the public 1 Ms observation. The 21 objects were classified, on the basis of deep VLT spectra, in two categories: 13 dusty star-forming galaxies showing [OII] emission, and 8 early-type galaxies with absorption features in their optical spectra. Only one emission line object has been individually detected; its very hard X-ray spectrum and the high intrinsic X-ray luminosity unambiguously reveal the presence of an obscured AGN. Stacking analysis of the remainder 12 emission line objects shows a significant detection with an average luminosity $L_X \approx 8 \times 10^{40}$ erg s^{−1} in the rest-frame 2–10 keV band. The stacked counts of the 8 passive galaxies do not provide a positive detection. We briefly discuss the implications of the present results for the estimate of the Star Formation Rate (SFR) in emission line EROs.

Subject headings: galaxies: active – galaxies: starburst – surveys – X-rays: galaxies

1. INTRODUCTION

Having the colors expected for high-redshift passive ellipticals, Extremely Red Objects (hereinafter EROs; $R - K > 5$) can be used as tracers of distant and old spheroids as a test for different cosmological models. The optical and near-infrared properties of EROs are also consistent with those expected for high-redshift dusty starbursts (e.g., HR 10, Cimatti et al. 1998; Smail et al. 1999) and Active Galactic Nuclei (AGN) reddened by dust (Pierre et al. 2001), for which the spectral energy distribution at these wavelengths is dominated by the host galaxy starlight (e.g., Mainieri et al. 2002).

The relative fraction of old ellipticals, dusty star-forming systems and obscured AGN among optically selected ERO samples is a key parameter in the study of the galaxy evolution and can constrain models which link the formation of massive elliptical galaxies and the onset of AGN activity (Almaini et al. 2002; Granato et al. 2001). A step forward in this direction has been recently achieved by the K20 survey (Cimatti et al. 2002a; Daddi et al. 2002). Deep VLT spectroscopy of a bright subsample at $K_s < 19.2$ has shown that EROs are nearly equally populated by old, passively evolving systems and dusty star-forming galaxies over a similar range of redshift ($z_{\text{mean}} \simeq 1$ for both the classes).

Sensitive hard X-ray observations provide a unique tool to uncover hidden AGN activity among the EROs population. Indeed, a sizeable fraction (10–30%) of hard X-ray sources recently discovered in deep *Chandra* and XMM surveys are associated with EROs and the fraction depends on the limiting fluxes reached both in the optical and X-ray bands (Alexander et al. 2002; Mainieri et al. 2002; Crawford et al. 2001). The hard X-ray spectra indicate that X-ray bright EROs are associated with high redshift, highly obscured AGNs.

Moreover, thanks to the very deep limiting fluxes reached by deep *Chandra* surveys (CDFs, Giacconi et al. 2002, hereinafter G02; CDFN, Brandt et al. 2001a) it is also possible to investigate via stacking analysis the average X-ray properties of fainter non-AGN sources (Brandt et al. 2001b; Hornschemeier et al. 2002; Nandra et al. 2002).

On the other hand, the AGN fraction in optically selected EROs samples is still very uncertain and possibly lower than 15% (Brusa et al. 2002; Alexander et al. 2002), suggesting that the bulk of the EROs population is not related to active phenomena. In fact, the majority of optically selected EROs remains undetected even at the deepest limiting fluxes of the 1 Ms *Chandra* Deep Field North observation (Alexander et al. 2002). Stacking analysis of those objects not individually detected suggests a softer X-ray spectrum than that of AGN-powered EROs and an X-ray to optical flux ratio consistent with that expected from normal elliptical galaxies, star-forming systems or low luminosity ($L_{2-10\text{keV}} < 10^{42}$ erg s^{−1}) nuclear activity at $z \simeq 1$.

Here we make use of the available spectroscopic information of a well-defined, magnitude limited ($K_s < 20$) subsample of EROs in the *Chandra* Deep Field–South area obtained as a part of the K20 survey (Cimatti et al. 2002a); the main aim of the present study is to investigate whether the X-ray emission properties of EROs are depending on the optical/infrared spectral properties.

Throughout the paper, a cosmology with $H_0 = 70$ km s^{−1} Mpc^{−1}, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$ is adopted.

2. THE SAMPLE

The ESO VLT K20 sample is one of the largest and most complete spectroscopic sample of galaxies with $K_s < 20$ available to date; the main scientific goal of this project is the comparison of

the observed redshift distribution of a complete sample of about 500 galaxies (87% complete to $K_s < 20$, 98% if photometric redshifts are included) with the predictions of galaxy formation models in order to obtain stringent clues on the formation and evolution of the present-day massive galaxies (Cimatti et al. 2002b). The K20 sample cover a sub-area (32.2 arcmin^2) of the *Chandra* Deep Field South (CDFS) and a 19.8 arcmin^2 field around the quasar 0055-2659.

Extremely Red Objects in the K20 sample were selected with the criterion of $R - K_s > 5$, and then classified on the basis of their optical spectra into two categories (Cimatti et al. 2002a): objects with an absorption line spectrum, consistent with that of early-type, passively evolving galaxies (hereinafter “old”) and objects showing [OII] $\lambda 3727$ emission line with an observed Equivalent Width $EW > 20$ Å over a red continuum, consistent with that of dusty star-forming systems (hereinafter “dusty”). The K20 EROs sample in the CDFS area includes 48 objects at $K_s < 20$. For the purposes of the present paper we consider only the 21 spectroscopically identified EROs which have been classified in the two categories described above: 13 “dusty” and 8 “old”, distributed on a similar range of redshifts, $z = 0.8 \div 1.6$.

3. X-RAY DATA ANALYSIS

The *Chandra* Deep Field South (CDFS) survey is a combination of 11 individual *Chandra* ACIS-I pointings observed in two years, from March 1999 to December 2000 (G02). The maximum effective exposure time is 942 ks, but it varies across the detector due to the CCD gaps and the rotation of the field-of-view in the different observations. In order to estimate the effective exposure time for each object, we have downloaded from the archive, along with the combined photon event list, all the events corresponding to the single pointings. We have calculated the exposure and instrument maps using standard CIAO 2.2.1 tools for each observation separately, and combined them to evaluate point-by-point the effective exposures. As the mirror vignetting is energy dependent, we calculated the exposure maps at two energies (1 keV and 4 keV), representative of the mean energy of the photons in the soft (0.5–2 keV) and hard (2–8 keV) bands.

At first, we have searched for nearby (within $2''$) X-ray sources found by WAVDETECT in the K20 region, with a false-positive probability threshold of 10^{-5} (Freeman et al. 2001). Only one source, a dusty ERO at $z = 1.327$, was individually detected; the optical position is almost coincident ($\Delta < 0.3''$, within the accuracy of the K-band position) with the *Chandra* source CXO CDFS J033213.9-274526 in the G02 catalog. The X-ray and optical properties of this object are discussed in Sect. 4.1.

In order to constrain average X-ray properties of the remaining individually undetected EROs, we have applied the stacking technique (Brandt et al. 2001b; Nandra et al. 2002) separately for the two classes of objects. The samples consist of 12 “dusty” and 8 “old” EROs, with average redshifts of $z = 1.053$ and $z = 1.145$, respectively.

For the photometry we used a circular aperture with a radius of $2''$ centered at the positions of the selected EROs. We have also checked different aperture radii ($1 - 4''$) and found that the

adopted one is an optimal extraction radius for the counts estimate, in agreement with the Nandra et al. (2002) findings. The counts were stacked in the standard soft, hard and full bands (0.5–2 keV, 2–8 keV, and 0.5–8 keV), and in the 1–5 keV band which roughly corresponds to the rest-frame 2–10 keV band at the mean redshift of the objects.

To verify whether the stacked net counts constitute a significant signal the local background has been estimated: in annular regions around each ERO centroid, as the average of eight circular regions (2 arcsec radius) for each ERO position and from a background map produced by the WAVDETECT algorithm. The different choices do not significantly affect the results in none of the adopted energy bands, except the 2–8 keV range where the results are strongly affected by the increased level of the instrumental background. Extensive Monte Carlo simulations (up to 100,000 trials) have been also carried out shuffling 12(8) random positions for “dusty” and “old” EROs respectively, using the same photometry aperture (2 arcsec). The random positions were chosen to lie in “local background regions” to reproduce the actual background as close as possible. The resulting distributions are well approximated by Gaussians. The corresponding detection confidence level is in all the cases in good agreement with that computed from Poisson statistic.

The results of the stacking analysis reported in Table 1 are referred to the background estimated as the average of the eight circular regions; in Fig. 1 we show the summed images in the full (0.5–8 keV) band. An excess of counts above the expected background level is clearly present for the 12 “dusty” EROs.

The signal is stronger in the full band (99.997% confidence level assuming a Poissonian distribution, corresponding to about 4.2σ) and it is still present, although at a slightly lower confidence level, in the soft and 1–5 keV band, while it is not statistically significant in the hard band. Assuming a $\Gamma = 2$ power law spectrum plus Galactic absorption ($N_H = 8 \times 10^{19} \text{ cm}^{-2}$), the 3σ upper limit on the average band ratio ($H/S < 0.85$, where H and S indicates the hard band and soft band counts, respectively), corresponds to an absorption column density lower than 10^{22} cm^{-2} (see Fig. 4 in Alexander et al. 2002).

Given that the X-ray spectrum of nearby starburst galaxies is well-represented by power-law models with slopes in the range $1.8 \div 2.5$ and low intrinsic absorption (e.g., Ptak et al. 1999; Ranalli et al. 2002a), we adopted $\Gamma = 2.1$ to compute X-ray fluxes and luminosities. With this assumption, the stacked count rate in the 1–5 keV band corresponds to an average 2–10 keV rest-frame luminosity of $\sim 8 \times 10^{40} \text{ erg s}^{-1}$, at the mean redshift $z = 1.053$.

The X-ray emission from “old” EROs remains undetected in all the considered energy bands (see Table 1 for details). Assuming a thermal emission model with $kT = 1 \text{ keV}$ and solar metallicity, fully consistent with the average X-ray spectrum of nearby elliptical galaxies (e.g., Pellegriini 1999), the 3σ upper limit on the 0.5–2 keV luminosity is $L_X < 10^{41} \text{ erg s}^{-1}$ (see also Vignali et al. 2002).




FIG. 1.— Stacked full-band images of the “dusty” (left panel) and “old” (right panel) EROs. The images are 9x9 arcsec and have been smoothed with a gaussian of $\sigma=1.5$ pixel (approx $0.7''$). The circles are centered on the stacking position and have a radius of $2''$. The detection significance of the summed counts is 4.2σ for the “dusty” and $< 2\sigma$ for the “old”. The bright spot in the right panel at about $4''$ from the stacked position, corresponds to an individually detected object.

TABLE 1
STACKING RESULTS

Class	band (keV)	counts total	counts BKG	c.l.	Exp. (Ms)	Flux [†]	$L_x^{\dagger}$
Dusty	0.5-8	172	124.2	99.997%	10.2	3.4 ± 1.3	20 ± 8
	0.5-2	67	37.9	99.995%	10.4	1.3 ± 0.45	7.4 ± 2.6
	1-5	83	61.4	99.6%	10.2	1.4 ± 0.5	8.0 ± 2.8
	2-8	105	86.3	97.2%	10.1	< 4.8	< 27
Old [‡]	0.5-8	102	92.1	$< 90\%$	7.0	< 7.5	< 40
	0.5-2	33	26.0	$< 90\%$	7.0	< 24	< 10

[†]: Fluxes (in units of 10^{-17} erg cm $^{-2}$ s $^{-1}$) and luminosity (in units of 10^{40} erg s $^{-1}$) are computed assuming a power-law spectrum with $\Gamma = 2.1$ and Galactic absorption for the “dusty” sample, and a thermal emission model with $kT=1$ keV and solar metallicity for the “old” sample. Errors are at the 90% confidence level.

[‡]: For the “old” sample we report only the results in the 0.5–8 keV and 0.5–2 keV bands. In the harder bands the results are always consistent, within 1σ , with the background fluctuations.

4. RESULTS

4.1. CXO CDFS J033213.9-274526: a type 2 AGN

The only ERO individually detected in the 1 Ms exposure is the highest redshift, highest luminous ($R=23.81$, $K_s=18.44$) object in the “dusty” sample. It is a very hard X-ray source ($F_{2-8\text{keV}}=7.34 \times 10^{-16}$ erg cm $^{-2}$ s $^{-1}$, $\Gamma = 1.4$ and $N_H=\text{Gal}$; G02) undetected in the soft band ($F_{0.5-2\text{keV}} < 5.48 \times 10^{-17}$ erg cm $^{-2}$). The observed band ratio ($H/S > 7.3$) implies an intrinsic X-ray column density larger than 4×10^{23} cm $^{-2}$, for a classic AGN power-law spectrum with $\Gamma = 1.8$). The observed 2–8 keV luminosity is $\sim 7 \times 10^{42}$ erg s $^{-1}$ which corresponds to an unabsorbed, rest-frame hard X-ray luminosity $\gtrsim 4 \times 10^{43}$ erg s $^{-1}$, typical of Seyfert galaxies. At the high-energy, therefore, CXO CDFS J033213.9-274526 is naturally classified as a Type 2, active object. In the optical domain it shows only a faint, but significant, [OII] $\lambda 3727$ emission feature. Unfortunately, at the expected [NeV] $\lambda 3426$ position a strong cosmic ray has affected the data. The lack of (broad) MgII $\lambda 2798$ emission indicates that obscured activity is taking place also in the optical nucleus.

¹ SFR (M $_{\odot}$ yr $^{-1}$) = $2.0 \times 10^{-40} L_{2-10\text{keV}}$ (erg s $^{-1}$)

Given the small area sampled and the spectroscopic completeness, the present ERO sample is not suitable for a reliable estimate of the AGN fraction among the EROs population. However, the present low detection rate of AGN among EROs (1/21) confirms that the bulk of the EROs population is not related to active phenomena.

4.2. Dusty EROs

The X-ray and optical data both suggest that starburst activity, rather than AGN, is powering “dusty” EROs. Indeed, following Cimatti et al. (2002a), we recomputed the average spectrum of the 12 EROs we used in the X-ray stacking analysis. The average optical spectrum is in excellent agreement with that published in Cimatti et al. 2002a (see their Fig. 2), derived from the complete K20 EROs sample, except for the presence of the AGN indicator [NeV] $\lambda 3426$; in the present “dusty” sample this feature can be safely ruled out ($EW < 1.5$ Å at the 3σ level), suggesting that the optical emission is dominated by star-forming systems.

The inferred, average X-ray luminosity is around one order of magnitude larger than that of normal spiral galaxies (Matsumoto et al. 1997) and it is similar to that of the starburst galaxy M82 (e.g., Griffiths et al. 2000; Kaaret et al. 2001); the average X-ray spectrum (see Sect. 3) is consistent with that measured for nearby star-forming galaxies (e.g. Dahlem, Weaver & Heckman 1998; Ptak et al. 1999).

We have also investigated whether the stacked signal could be due to a few individual sources just below the detection threshold. This is particularly relevant for our sample which may contain a few sub-threshold AGN responsible for the entire stacked signal. Five out of 12 “dusty” EROs are individually “detected” at the 95–99% confidence level. The X-ray luminosity for these “marginally detected” EROs never exceeds 6×10^{41} erg s $^{-1}$ in the full-band excluding the presence of a bright AGN among the stacked dusty EROs. Although the presence of a low-luminosity AGN (LLAGN) can not be completely ruled out on the basis of the present analysis, in the following we assume that the “dusty” EROs are mainly powered by starburst related processes.

4.2.1. Star Formation Rate: the hard X-rays view

There is an increasing evidence that the hard X-ray luminosity of star-forming objects can be used as a star formation rate (SFR) indicator (Ranalli, Comastri, & Setti 2002a,b; Nandra et al. 2002; Bauer et al. 2002).

Using the Ranalli et al. relation¹ we have derived an average SFR for the EROs in the “dusty” sample of $5\text{--}44$ M $_{\odot}$ yr $^{-1}$, taking into account the observed dispersion between the far-infrared and X-ray luminosities quoted by Ranalli et al., and the uncertainties on the hard X-ray luminosity. The average SFR derived from the the observed [OII] luminosity (Kennicutt 1998) is $\text{SFR}([\text{OII}])_{\text{obs}}=2.7$ M $_{\odot}$ yr $^{-1}$. The discrepancy between the optical and X-ray SFR estimates confirms that the optical emission lines are severely obscured by dust and

the optical value can be addressed as a *lower limit*. In order to provide a dereddened SFR estimate, the average spectrum was compared with template spectra of star forming galaxies and dusty systems (as extensively discussed in Cimatti et al. 2002a); the best-fit estimates for the reddening are in the range $E(B-V) = 0.5 \div 0.8$, which implies an intrinsic average SFR of the order $37\text{--}110\text{ M}_\odot\text{ yr}^{-1}$ according to the Kennicutt (1998) relation. Although the two estimates suffer of large uncertainties, mainly related to the scatter of the various SFR indicators, they can be reconciled if the lowest extinction correction (e.g., $E(B-V) \sim 0.5$, or even lower) is applied to correct the [OII] luminosity.

From the derived X-ray SFR it is possible to estimate the average far-infrared (60–100 μm) and radio (1.4 GHz) emission of the objects in our sample: $L_{\text{FIR}} = 10^{10.4} \div 10^{11.4}\text{ L}_\odot$; $L_{1.4\text{GHz}} = 2 \div 17.6 \times 10^{29}\text{ erg s}^{-1}\text{ Hz}^{-1}$ (Kennicutt 1998). The relatively low value of L_{FIR} would naturally explain the low detection rate of EROs in submillimeter continuum follow-up observations, typically sensitive to detect Ultra-Luminous Infrared Galaxies (ULIGs, $L > 10^{12}\text{ L}_\odot$) at $z \gtrsim 1$ (e.g., Mohan et al. 2002). The “dusty” ERO population would then be representative of less luminous dusty star-forming systems, with respect to submillimeter-selected high- z galaxies, as already pointed out in Cimatti et al (2002a).

The inferred radio luminosity corresponds to K-corrected ($S_\nu \propto \nu^{-\alpha}$, $\alpha = 0.7$) radio fluxes in the range $4\text{--}38\mu\text{Jy}$. The lack of deep radio observations in the *Chandra* Deep Field South area prevents to test these predictions; however, it is interesting to note that very deep 1.4 GHz VLA observations of a sample of optically selected EROs (Smail et al. 2002) agree with our estimates (about 85% of the EROs in their sample is not detected in the radio band at fluxes higher than $\sim 40\mu\text{Jy}$).

5. SUMMARY

Thanks to the deep limiting fluxes reached in the 1 Ms exposure of the *Chandra* Deep Field South and to the subarcsec positional accuracy of the X-ray satellite, it was possible to constrain via stacking analysis the high-energy properties of a well-defined, spectroscopically identified sample of EROs, belonging to the K20 survey.

The dichotomy in the optical classifications of EROs seems to be present also in the X-ray band: “dusty” objects are relatively bright X-ray sources, while “old” EROs are below the detection threshold.

The only object (CXO CDFS J033213.9-274526) individually detected by *Chandra* is a luminous, highly absorbed AGN ($N_H > 4 \times 10^{23}\text{ cm}^{-2}$) at $z = 1.327$.

Using the hard X-ray luminosity as a SFR indicator, which is essentially free from intrinsic absorption, we have estimated a relatively low SFR ($5\text{--}44\text{ M}_\odot\text{ yr}^{-1}$) with respect to the estimate based on the reddening-dependent [OII] emission; our estimate is consistent with the results of recent far-infrared and radio observations.

Deep X-ray observations of large and complete samples of spectroscopically identified EROs could provide a powerful tool to estimate, via stacking analysis, the average star formation rate in “dusty” systems and to better assess the AGN fraction among the ERO population.

The authors acknowledge partial support by ASI I/R/103/00 and I/R/113/01 contracts and MIUR Cofin-00-02-36 grant. MB and AC thank P. Ranalli, C. Vignali and G. Zamorani for useful comments and discussions. We thank the anonymous referee for his/her helpful comments.

REFERENCES

- Alexander, D.M., Vignali, C., Bauer, F.E., Brandt, W.N., Hornschemeier, A.E., Garmire, G.P., & Schneider, D.P. 2002, *AJ*, 123, 1149
 Almaini O., et al. 2002, *MNRAS*, in press (astro-ph/0108400)
 Bauer, F.E., Alexander, D.M., Hornschemeier, A.E., Vignali, C., Garmire, G.P., & Schneider, D.P. 2002, *AJ* in press (astro-ph/0207433)
 Brandt, W.N., et al. 2001a, *AJ*, 122, 2810
 Brandt, W.N., Hornschemeier, A.E., Schneider, D.P., Alexander, D.M., Bauer, F.E., Garmire, G.P., & Vignali, C. 2001b, *ApJ*, 558, L5
 Brusa, M., Comastri, A., Daddi, E., Cimatti, A., & Vignali, C. 2002, in “X-ray spectroscopy of AGN with *Chandra* and XMM-Newton”, eds. Boller T., Komossa S., Kahn S., Kunieda H. (astro-ph/0204251)
 Cimatti, A., Andreani, P., Röttgering, H., & Tilanus, R. 1998, *Nature*, 392, 895
 Cimatti, A., et al. 2002a, *A&A*, 381, L68
 Cimatti, A., et al. 2002b, *A&A*, 391, L1
 Cimatti, A., et al. 2002c, *A&A*, 392, 395
 Crawford, C.S., Fabian, A.C., Gandhi, P., Wilman, R.J., & Johnstone, R.M. 2001, *MNRAS*, 324, 427
 Daddi, E., et al. 2002, *A&A*, 384, L1
 Dahlem, M., Weaver, K.A., & Heckman, T.M. 1998, *ApJS*, 118, 401
 Elston, R., Rieke, G.H., & Rieke, M.J. 1988, *ApJ*, 331, L77
 Freeman, P.E., Kashyap, V., Rosner, R. & Lamb, D. 2002, *ApJS*, 138, 185
 Giacconi, R., et al. 2002, *ApJS*, 139, 369 (G02)
 Granato, G.L., Silva, L., Monaco, P., Panuzzo, P., Salucci, P., De Zotti, G., & Danese, L. 2001, *MNRAS*, 324, 757
 Griffiths, R.G., et al. 2000, *Science*, 290, 1325 (astro-ph/0205371)
 Hornschemeier, A.E., Brandt, W.N., Alexander, D.M., Bauer, F.E., Garmire, G.P., Schneider, D.P., Bautz, M.W., & Chartas, G. 2002, *ApJ*, 568, 82
 Kaaret, P., Prestwich, A.H., Zezas, A., Murray, S.S., Kim, D.W., Kilgard, R.E., Schlegel, E.M., & Ward, M.J. 2001, *MNRAS*, 321, L29
 Kennicutt, R.C. 1998, *ARA&A*, 36, 189
 Mainieri, V., Bergeron, J., Hasinger, G., Lehmann, I., Rosati, P., Schmidt, M., Szokoly, G., & Della Ceca, R. 2002, *A&A* 393, 425
 Matsumoto, H., Koyama, K., Awaki, H., Tsuru, T., Loewenstein, M., Matsushita, K. 1997, *ApJ*, 482, 133
 Mohan, N. R., Cimatti, A., Röttgering, H.J.A., Andreani, P., Severgnini, P., Tilanus, R.P.J., Carilli, C.L., & Stanford, S.A. 2002, *A&A*, 383, 440–456
 Nandra, K., Mushotzky, R.F., Arnaud, K., Steidel, C.C., Adelberger, K.L., Gardner, J.P., Teplitz, H.I., & Windhorst, R.A. 2002, *ApJ*, 576, 625
 Pellegrini, S. 1999, *A&A*, 351, 487
 Ptak, A., Serlemitsos, P., Yaqoob, T., & Mushotzky, R.F. 1999, *ApJS*, 120, 179
 Pierre M., et al. 2001, *A&A*, 372, L45
 Ranalli, P., Comastri, A., Setti, G.C. 2002a, *A&A*, in press
 Ranalli, P., Comastri, A., & Setti, G.C. 2002b, in “New Visions of the X-ray Universe in the XMM-Newton and Chandra Era”, ed. F. Jansen (astro-ph/0202241)
 Rosati, P., et al. 2002, *ApJ*, 566, 667
 Smail, I., Ivison, R.J., Kneib, J.P., Cowie, L.L., Blain, A.W., Barger, A.J., Owen, F.N., & Morrison, G., 1999, *MNRAS* 308, 1061
 Smail, I., Owen, F.N., Morrison, G.E., Keel, W.C., Ivison, R.J., & Ledlow, M.J. 2002, *ApJ* in press (astro-ph/0208434)
 Vignali, C., Alexander, D.M., Bauer, F.E., Brandt, W.N., Hornschemeier, A.E., Garmire, G.P., & Schneider, D.P., proceedings of the 5th Italian AGN Meeting “Inflows, Outflows and Reprocessing around black holes” (2002), astro-ph/0209415